



TECHNICAL INFORMATION BULLETIN
FOR THE USE OF REGALIA® SC TO CONTROL DOWNY MILDEW
ON LEAFY VEGETABLE CROPS

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF CALIFORNIA AND ARIZONA.

REGALIA® SC
EPA Reg. No. 84059-2

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROP	DISEASE	RATE	USE RECOMMENDATIONS
Leafy Vegetable Crops such as Lettuce, Celery, Spinach, Parsley, and Radicchio	Downy Mildew (<i>Bremia lactucae</i> , <i>Peronospora spp.</i>)	0.5% v/v in 50 gallons of water (1 quart in 50 gallons of water).	For control of Downy Mildew, apply Regalia® SC preventively at 0.5% v/v in 50 gallons of water (1 quart in 50 gallons of water), adding a non-ionic surfactant according to manufacturer label directions. Repeat applications at 7- to 14-day intervals. Regalia® SC may be applied up to and including the day of harvest (0-day PHI).

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

Regalia® SC is a registered trademark of Marrone Bio Innovations, Inc.

© Copyright Marrone Bio Innovations, Inc., 2009

Marrone Bio Innovations, Inc.
2121 Second Street, Suite B-107
Davis, California 95618
Telephone (530) 750-2800
www.marronebioinnovations.com