



TECHNICAL INFORMATION BULLETIN
FOR THE USE OF REGALIA®
FOR CONTROL OF
ALTERNARIA BROWN SPOT
ON CITRUS

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE RECOGNIZED.

REGALIA®
EPA Reg. No. 84059-3

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROP	DISEASE	RATE	USE RECOMMENDATIONS
Citrus crops including Orange, Grapefruit, Lemon, Tangerine, Tangelo, and Pummelo.	Alternaria Brown Spot (<i>Alternaria alternata</i>)	Apply Regalia® at a rate of 2-4 qts per acre when applied alone or 1-4 qts per acre when tank mixed with another fungicide in dilute applications to the point of runoff. Apply Regalia® preventatively in 50-100 gallons of water per acre.	For improved performance, use Regalia in a tank mix or rotational program with other registered fungicides. Make first application at spring flush, second application at petal fall and thereafter according to predictive weather-based models until fruit is no longer susceptible to the disease.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

Regalia® is a registered trademark of Marrone Bio Innovations, Inc.
© Copyright Marrone Bio Innovations, Inc., 2010

Marrone Bio Innovations, Inc.
2121 Second Street, Suite B-107
Davis, California 95618
Telephone (530) 750-2800
www.marronebioinnovations.com