



TECHNICAL INFORMATION BULLETIN
 FOR THE USE OF MAJESTENE™ BIONEMATICIDE
 ON BLACKBERRIES AND RASPBERRIES

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(Ee) RECOMMENDATIONS ARE RECOGNIZED.

MAJESTENE™
EPA Reg. No. 84059-14

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

| Crop | Target Nematodes | Application Method | Product Use Rate per Application | Application Instructions |
|--|---|----------------------------|--|---|
| Blackberries and raspberries | Suppression of plant parasitic nematodes including root lesion and dagger nematodes | Post-planting applications | 2 gallons per acre as a ground spray 2 gallons per acre in a pressurize drip or sprinkler irrigation system | Apply Majestene as a 2-4 foot band application over the row in at least 50 gallons of water per acre prior to imminent rainfall or followed by overhead irrigation of 0.25 – 0.5 inches of water to move the product into the top 2-3 inches of soil. Make the initial application when the soil temperature is at least 45°F at an 8 inch depth. Additional applications can be made on a 3-4 week interval prior to imminent rainfall or followed by sprinkler irrigation. Inject Majestene mid-way into the irrigation event to allow for incorporation of the product into the soil profile. Additional applications can be made at 21-day or greater intervals. |
| MAJESTENE™ has a pre-harvest interval (PHI) of 0 days . Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours . | | | | |

Nematode management considerations - To obtain maximum results in a nematode management program, it is important to take soil samples prior to application to assess nematode populations and determination of an appropriate management strategy. Utilization of the services of a field person or crop consultant who is knowledgeable in nematode management is advised.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

MAJESTENE™ is a trademark of Marrone Bio Innovations, Inc.
 © Copyright Marrone Bio Innovations, Inc., 2014
 Marrone Bio Innovations, Inc. www.marronebioinnovations.com