



TECHNICAL INFORMATION BULLETIN
FOR THE USE OF MAJESTENE® BIONEMATICIDE
New Tree Plantings

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EЕ) RECOMMENDATIONS ARE RECOGNIZED.

MAJESTENE®
EPA Reg. No. 84059-14

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop	Target Nematodes	Application Method	Product Use Rate per Application	Application Instructions
New Tree /Orchard Plantings (includes Vine, Fruiting and Nut Trees, and Ornamental Trees)	For control or suppression of plant parasitic nematodes including: Dagger, Lesion, Reniform, Ring, Root- knot, Cyst, Sting, Stubby-root, and Stunt, nematodes.	Post-planting application	1-4 gallons per acre	Apply the desired rate by chemigation into the root zone through low-pressure microsprinkler, drip, or trickle irrigation equipment. For optimal results soil should be pre-wetted prior to chemigation. Multiple applications may be necessary to achieve desired results. An application interval of 21-35 days is recommended. Inject Majestene® in the final 30-60 minutes of an irrigation cycle, sufficient to wet the root zone, maintaining a minimum concentration of 0.05% v/v to optimize product performance.
		Re-plant applications	1-4% v/v	For situations where individual trees are being replanted, saturate the root ball and the soil at the planting site to the depth/ volume of the anticipated root zone of the new planting with a 1-4% v/v solution of Majestene®. Subsequent applications should follow within 28 days.

MAJESTENE® has a **pre-harvest interval (PHI)** of **0 days**.
Do not enter or allow worker entry into treated areas during the **restricted-entry interval (REI)** of **4 hours**.

MAJESTENE® has no plant back or rotational restrictions and the active ingredient is tolerance exempt – No Maximum Residue Levels (MRL)



Nematode management considerations - To obtain maximum results in a nematode management program, it is important to take soil samples prior to planting to assess nematode populations and determine an appropriate management strategy. Utilization of the services of a field person or crop consultant who is knowledgeable in nematode management is advised.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

MAJESTENE® is a registered trademark of Marrone Bio Innovations, Inc.

© Copyright Marrone Bio Innovations, Inc., 2017

Marrone Bio Innovations, Inc.

1540 Drew Ave.

Davis, California 95618

Telephone (530) 750-2800

www.marronebioinnovations.com